

Prescription Treatment[®] brand Sorex

Pellet Place Pack

- Kills Rats & Mice
- Ready to use
- For indoor and outdoor use.

ACTIVE INGREDIENT:

Difenacoum

OTHER INGREDIENTS:

BY WT.

0.005%

99.995%

TOTAL: 100.000%

EPA Reg. No. 47629-17-499

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Keep away from children, domestic animals and pets. All handlers (including applicators) must wear shoes plus socks, and gloves. Any person who retrieves carcasses or unused bait following application of this product must wear waterproof gloves.

USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash hands thoroughly after applying bait and before eating, drinking, chewing gum, using tobacco or using the toilet and change into clean clothing.

FIRST AID

HAVE LABEL WHEN OBTAINING TREATMENT ADVICE.

IF SWALLOWED: Call a poison control center, doctor or 1-800-832-HELP immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor.

TREATMENT FOR PET POISONING

If animal eats bait, call veterinarian at once.

NOTE TO PHYSICIAN OR VETERINARIAN

Anticoagulant Difenacoum: If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. **For humans or dogs** that have ingested this product and/or have obvious poisoning symptoms (bleeding or prolonged prothrombin times), give Vitamin K₁ intramuscularly or orally. **For dogs**, check prothrombin times every 3 days until values return to normal (up to 30 days). Blood transfusions may be needed.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and wildlife. Dogs and predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten bait. Do not apply this product directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

READ THIS LABEL! Read this entire label and follow all use directions and use precautions. Use only for the sites, pests, and application methods described on this label.

IMPORTANT: Do not expose children, pets, or nontarget animals to rodenticides. To help prevent accidents:

1. Store unused product out of reach of children and pets.
2. Apply product in bait stations for all outdoor, above ground placements.

3. Use tamper-resistant bait stations for all placements where children, pets, nontarget mammals or birds may access bait. These stations must be resistant to destruction by dogs and by children under 6 yrs of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from bait stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
4. Dispose of product container and unused, spoiled, or unconsumed bait as specified on this label.

USE RESTRICTIONS

This product may be used to control Norway rats, roof rats, and house mice only at the **Urban** and **Non-urban** sites identified below. **Do not apply further than 50 ft from buildings.** Do not use this product in sewers. Do not place bait near to or inside of ventilation duct openings. Do not contaminate water, food, feed, food or feed handling equipment, milk or meat handling equipment, or any area where there is a possibility of contaminating food, feed, milk, or meat, or surfaces that come into contact with food, feed, milk, or meat. Do not use in edible product areas of food or feed processing plants or other areas where food or feed is commercially prepared or processed. All outdoor, above ground placements of this product must be made in bait stations. Tamper-resistant bait stations must

Sorex Pellet Place Pack

be used for outdoor and indoor placements of this product if children, pets, non-target mammals, or birds may access bait. All placements of this product in USDA-inspected facilities must be in tamper-resistant bait stations. Do not broadcast bait. Wear water-resistant gloves when handling bait.

Do not sell this product in individual containers holding less than 16 lbs of bait.

Urban Areas: This product may only be used in and around homes, industrial, commercial, and public buildings, and similar man-made structures; in transport vehicles (ships, trains, and aircraft) and in and around related port or terminal buildings; and in alleys.

Non-Urban Areas: This product may only be used in and around homes, and agricultural buildings.

SELECTION OF TREATMENT AREAS

Determine areas where rats and mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, beside burrows, in corners and concealed places, between floors and walls, or in locations where rats, mice, or their signs have been seen. Protect bait from rain and snow. Remove as much alternative food as possible.

APPLICATION DIRECTIONS

Norway Rats and Roof Rats: Use 3 - 10 place packs per placement, using bait stations as required under **"IMPORTANT"**. Space placements at intervals of 15 - 30 ft in infested areas pertaining to structures. Maintain an uninterrupted supply of fresh bait for at least 10 days or until fresh signs of feeding by rats cease to appear.

House Mice: Open place pack and apply 0.25 - 0.5 oz of bait per placement, using bait stations as required under **"IMPORTANT"**. Space placements at intervals of 8 - 12 ft in infested areas pertaining to structures. Larger quantities (up to 1 unopened 1.5 oz place pack may be placed in areas of extremely high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days or until fresh signs of feeding by mice cease to appear.

Follow-up: Replace contaminated or spoiled bait immediately. Using water-resistant gloves, collect and properly dispose of dead animals and leftover bait, as well as any bait that has become accessible to non-target animals. To discourage reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. For continuous sources of infestation, set up permanent bait stations and replenish them as needed.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store only in original container in a cool, dry place inaccessible to children and pets. Keep containers closed and away from other chemicals.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container.

Triple rinse (or equivalent) then offer for recycling or reconditioning; or puncture and dispose of in a sanitary landfill; or incineration.

NOTICE: Buyer assumes all risks of use, storage or handling of this product not in accordance with label directions.

Note: Seller warrants that this product complies with the specifications expressed in this label. To the extent consistent with applicable law, Seller makes no other warranties, and disclaims all other warranties, express or implied, including but not limited to warranties of merchantability and fitness for the intended purpose. To the extent consistent with applicable law, Seller's liability or default, breach or failure under this label shall be limited to the amount of the purchase price. To the extent consistent with applicable law, Seller shall have no liability for consequential damages.

Manufactured for:
Whitmire Micro-Gen Research Laboratories, Inc.®
3568 Tree Court Industrial Blvd.
St. Louis MO 63122-6682
Questions? Call 1-800-777-8570
© 2011 Whitmire Micro-Gen Research Laboratories, Inc.®

NOTE: This specimen label is for informational purposes only. All uses may not be approved in all states. See labeling which accompanied product for Directions for Use or call 800-777-8570 for more information.

120102-30